



# BOTANISKA NOTISER

1971, VOL. 124

ISSUED BY  
THE LUND BOTANICAL SOCIETY

EDITED BY  
*ARNE STRID*

*Established in 1839*

DISTRIBUTOR: C. W. K. GLEERUP, LUND, SWEDEN

#### Distribution Dates

Fasc. 1, pp. 1—178, on 1 April 1971  
Fasc. 2, pp. 179—310, on 9 July 1971  
Fasc. 3, pp. 311—432, on 29 Oct. 1971  
Fasc. 4, pp. 433—522, on 30 Dec. 1971

# Contents

ADATIA, R. D., SHARMA, Y. B. and VIJAYA-RAGHAVAN, M. R.: Studies in the Gesneriaceae I. Morphology and Embryology of <i>Platystemma violoides</i> Wall. . . . .	25	MOORE, D. M.: <i>Tapeinia</i> Comm. ex Juss. (Iridaceae) and the Rediscovery of <i>Galaxia obscura</i> Cav. . . . .	82
AREKAL, D., RAJESHWARI, S. and SWAMY, S. N. R.: Contribution to the Embryology of <i>Scoparia dulcis</i> L. . . . .	237	— and CHATER, A. O.: Studies of Bipolar Disjunct Species I. <i>Carex</i> . . . . .	317
BENTZER, B., BOTHMER, R. von, ENGSTRAND, L., GUSTAFSSON, M. and SNOGERUP, S.: Some Sources of Error in the Determination of Arm Ratios of Chromosomes . . . . .	65	MUNTING, A.: Göte Turesson in Memoriam . . . . .	419
BOTHMER, R. von, ENGSTRAND, L., GUSTAFSSON, M., PERSSON, J., SNOGERUP, S. and BENTZER, B.: Clonal Variation in Populations of <i>Anemone nemorosa</i> L. . . . .	505	NILSSON, Ö.: Studies in Montia L., Claytonia L. and Allied Genera V. The Genus <i>Montiastrum</i> (Gray) Rydb. . . . .	87
BRAMWELL, D., HUMPHRIES, C. J., MURRAY, B. G. and OWENS, S. J.: Chromosome Numbers in Plants from the Canary Islands . . . . .	376	— Studies in Montia L., Claytonia L. and Allied Genera VI. The Genera Limnalsine Rudh. and Maxia Ö. Nilss. . . . .	187
DAHLGREN, R.: Current Topics. Multiple Similarity of Leaf between Two Genera of Cape Plants, <i>Cliffortia</i> L. (Rosaceae) and <i>Aspalathus</i> L. (Fabaceae) . . . . .	292	— and LASSEN, P.: Chromosome Numbers of Vascular Plants from Austria, Mallorca and Yugoslavia . . . . .	270
— Chromosome Numbers in the South African Genus <i>Aspalathus</i> L. (Fabaceae) . . . . .	383	— and SNOGERUP, S.: Drawings of Scandinavian Plants 45—49. <i>Juncus</i> L. . . . .	1
— KARLSSON, T. and LASSEN, P.: Studies on the Flora of the Balearic Islands I. Chromosome Numbers in Balearic Angiosperms . . . . .	249	— Ditto 50—54. <i>Juncus</i> L. . . . .	179
— and RAO, V. S.: The Genus <i>Oftia</i> Adans. and its Systematic Position . . . . .	451	— Ditto 55—59. <i>Juncus</i> L. . . . .	311
HAINES, R. W. and LYÉ, K. A.: Studies in African Cyperaceae IV. <i>Lipocephala</i> R. Br., <i>Hemicarpha</i> Nees, and <i>Isolepis</i> R. Br. . . . .	473	— Ditto 60—64. <i>Juncus</i> L. . . . .	435
HALE, M. E., Jr.: Studies on <i>Parmelia</i> Subgenus <i>Xanthoparmelia</i> (Lichenes) in South Africa . . . . .	343	NORDENSTAM, B.: New South African Compositae . . . . .	9
HAWKSWORTH, D. L.: Regional Studies in <i>Alectoria</i> (Lichenes) I. The Central and Southern African Species . . . . .	122	— Cytogeography of the Genus <i>Hedypnois</i> (Compositae) . . . . .	483
HOLMQUIST, C.: Northerly Localities for three Aquatic Plants, <i>Lemna trisulca</i> L., <i>Ceratophyllum demersum</i> L., and <i>Myriophyllum spicatum</i> L. . . . .	335	PERSSON, J.: Studies in the Aegean Flora XIX. Notes on <i>Alyssum</i> and Some Other Genera of Cruciferae . . . . .	399
KERS, L. E.: Notes on the Flora of Angola 16		RAO, V. S.: The Disk and its Vasculature in the Flowers of Some Dicotyledons . . . . .	442
— <i>Monsonia parvifolia</i> Schinz (Geraniaceae), a Species with Concealed Spurs . . . . .	208	RECHINGER, K. H.: <i>Potentilla Goulandrii</i> , <i>Dianthus Stamatiadae</i> , <i>Onosma psamophilum</i> , drei neue Arten der griechischen Flora . . . . .	75
LYÉ, K. A.: Studies in African Cyperaceae II. The Genus <i>Oxycaryum</i> Nees . . . . .	280	— <i>Lithospermum Goulandriorum</i> Rech. f., eine neue, bemerkenswerte Art der griechischen Flora . . . . .	355
— Studies in African Cyperaceae III. A New Species of <i>Schoenoplectus</i> and some New Combinations . . . . .	287	RUNGBY, S.: A Contribution to the Bryophytic Flora of Attica and Peloponnes . . . . .	228
		SAHLIN, C. I.: Zur Taraxacum-Flora Niedersachsens . . . . .	497
		SNOGERUP, S.: New Species of <i>Bupleurum</i> from Turkey . . . . .	359
		STRID, A.: Past and Present Distribution of <i>Nigella arvensis</i> L. ssp. <i>arvensis</i> in Europe . . . . .	231
		— Chromosome Numbers in some Albanian Angiosperms . . . . .	490
		SUNESON, S.: A Bifurcate Frond of <i>Laminaria digitata</i> from Sweden . . . . .	277
		VENKATA RAO, C.: Anatomy of the Inflorescence of some Euphorbiaceae. With a Discussion on the Phylogeny and Evolution of the Inflorescence including the Cyathium . . . . .	39

WEIMARCK, G.: Variation and Taxonomy of <i>Hierochloë</i> (Gramineae) in the Northern Hemisphere .....	129
WIGH, K. and STRANDHEDÉ, S.-O.: Chromosome Numbers in Some Swedish and Danish Mosses .....	213
 Reviews of Botanical Literature	
CULBERSON, C. F.: Chemical and Botanical Guide in Lichen Products. (By O. ALMBORN) .....	430
— Ditto. Supplement. (By O. ALMBORN)	430
DAVIS, P. H., HARPER, P. C. and HEDGE, I. C. (eds.): Plant Life of South-West Asia. (By B. NORDENSTAM) .....	429
FOTT, B.: Algenkunde Ed. 2. (By T. v. WACHENFELDT) .....	520
FREY, E.: Flechten. Unbekannte Pflanzenwelt. (By O. ALMBORN) .....	522
GUILLARMOD, J.: Flora of Lesotho (Basutoland). (By B. NORDENSTAM) .....	426
HALE, M. E.: The Lichens. (By O. ALMBORN) .....	431
INGOLD, C. T.: Fungal Spores. Their Liberation and Dispersal. (By J. ERIKSSON)	425
KUNKEL, G.: Helechos cultivados. (By O. ALMBORN) .....	308
— Arboles Exóticos. Los árboles cultivados en Gran Canaria I. (By O. ALMBORN) .....	309
NYHOLM, E.: Illustrated Moss Flora of Fennoscandia 2 (6). (By E. VON KRUSSENSTJERNA and B. LANGE) .....	305
PANKOW, H.: Algenflora der Ostsee. I. Benthos (Blau-, Grün-, Braun- und Rotalgen. (By B. HÄGERHÄLL) .....	520
POELT, J.: Bestimmungsschlüssel europäischer Flechten. (By O. ALMBORN) .....	521
PRESTON, R. D. (ed.): Advances in Botanical Research. 3. (By the Editor) .....	307
RECHINGER, K. H.: Flora Iranica. Fasc. 58—89. (By B. NORDENSTAM) .....	427
SAVAGE, J. M.: Evolution. Ed. 2. (By B. BENTZER) .....	429
SCHUSTER, R. M.: The Hepaticae and Anthocerotae of North America East of the Hundredth Meridian 1—2. (By O. ALMBORN) .....	176
STEBBINS, G. L.: Chromosomal Evolution in Higher Plants. (By the Editor) .....	307
TÉTÉNYI, P.: Infraspecific Chemical Taxa of Medicinal Plants. (By U. OLSSON) ..	176

# Index

The index covers only taxa which have been treated in more detail. New taxa and new combinations are printed in boldface. \* indicates illustration(s) (also of chromosomes) and m refers to a distribution map.

<i>Abietinella abietina</i>	217*	<i>Arum italicum</i>	250*
<i>Acalypha indica</i>	41 f*, 56*	<i>Aspalathus</i>	292 ff*, 383 ff
— sp.	56*	— <i>abietina</i>	384*, 396
<i>Aganosma cymosa</i>	447*	— <i>aculeata</i>	384*
<i>Ajuga iva</i>	257*	— <i>arida</i>	396
<i>Albizia lebbek</i>	447*	— — <i>ssp. arida</i>	385*
<i>Alectoria</i>	122 ff	— <i>aspalathoides</i>	385*
— <i>angolensis</i>	123	— <i>asparagooides</i> ssp. <i>asparagooides</i>	385*
— <i>bicolor</i>	127	— <i>calcarata</i>	385*
— cf. <i>chalybeiformis</i>	123	— <i>costulata</i>	386*
— <i>chalybeiformis</i> f. <i>terrestris</i>	127	— <i>decora</i>	386*
— <i>epichrysa</i>	127	— <i>desertorum</i>	386*
— <i>fuscescens</i> var. <i>nigrescentifusca</i>	123 f	— <i>dianthopora</i>	386*
— <i>jubata</i>	127	— <i>excelsa</i>	387*
— — var. <i>prolixa</i>	127	— <i>filicaulis</i>	387*
— <i>motykae</i>	124*	— <i>hispida</i> ssp. <i>hispida</i>	388*
— <i>ruwenzoriensis</i>	124 f*	— <i>macrocarpa</i>	389*
— <i>smithii</i> f. <i>esorediata</i>	126	— <i>marginata</i>	389*
— <i>sulcata</i>	127	— <i>pinguis</i> ssp. <i>australis</i>	390*
<i>Alkanna lutea</i>	252*	— <i>quinquefolia</i> ssp. <i>virgata</i>	390*
<i>Allium subhirsutum</i>	257*	— <i>rugosa</i> ssp. <i>rugosa</i>	390*
<i>Althaea hirsuta</i>	259*	— <i>sericea</i> ssp. <i>sericea</i>	391*
<i>Alyssoides cretica</i>	399*	— <i>spinosa</i> ssp. <i>spinosa</i>	391*
— <i>utriculata</i>	400	— <i>tridentata</i> ssp. <i>rotunda</i>	392*
<i>Alyssum</i>	400 ff	— <i>tylodes</i>	392*
— <i>alyssoides</i>	406*	— <i>uniflora</i> ssp. <i>uniflora</i>	392*, 394
— <i>foliosum</i>	400 f* m	— — <i>ssp. willdenowiana</i>	392*
— <i>fulvescens</i>	401 f*	— <i>villosa</i>	392*
— <i>hirsutum</i>	402*	— <i>vulpina</i>	392*
— <i>minus complex</i>	402 ff	— <i>zeyheri</i>	392*
— <i>minus</i>	403 f*	<i>Asteriscus aquaticus</i>	250*
— <i>siculum</i>	406 ff* m	<i>Astragalus epiglottis</i> ssp. <i>epiglottis</i>	254*
— <i>smyrnaeum</i>	404*	<i>Asterolinum linum-stellatum</i>	260*
— <i>strigosum</i>	404 ff	<i>Asystasia gangetica</i>	443*
— — <i>ssp. cedrorum</i>	404 ff* m	<i>Atrichium undulatum</i>	223*
— — <i>ssp. strigosum</i>	404 ff*	<i>Aurinia saxatilis</i>	411 ff
— <i>umbellatum</i>	406*	— — <i>ssp. megalocarpa</i>	411 ff* m
<i>Androcymbium psammophilum</i>	377*	— — <i>ssp. orientalis</i>	411 ff* m
<i>Andrographis paniculata</i>	446*	— — <i>ssp. saxatilis</i>	411 ff* m
<i>Anemone nemorosa</i>	505 ff*	<i>Brachythecium albicans</i>	219*
<i>Anthostema senegalensis</i>	58*	— <i>plumosum</i>	219*
<i>Argyranthemum anethifolium</i>	380*	— <i>populeum</i>	219 f*
— <i>callichrysum</i>	380*	— <i>reflexum</i>	220*
— <i>canariense</i>	380*	— <i>rivulare</i>	220*
— <i>foeniculaceum</i>	379*	— <i>rutabulum</i>	220*
— <i>frutescens</i> var. <i>erithmifolium</i>	379 f*	— <i>velutinum</i>	220 f*
— — var. <i>frutescens</i>	379*	<i>Brassica balearica</i>	252*
— — var. <i>gracilescens</i>	380*	<i>Breynia</i>	56*
— <i>grandiflorum</i>	380*	— <i>rhamnooides</i>	56*
— <i>haouarytheum</i>	380*	<i>Briza maxima</i>	376 f*
<i>Arisarum vulgare</i>	250*	— <i>media</i>	377*
<i>Aristida adscensionis</i>	20*	<i>Bryum pseudotriquetrum</i>	214 f*

Bupleurum . . . . .	359 ff	Geranium ocellatum . . . . .	20 f
— apiculatum . . . . .	362 m	Gonospermum gomerae . . . . .	381*
— davisii . . . . .	362 ff*	Haplophragma adenophyllum . . . . .	443*
— eginense . . . . .	368 f*	Hedypnois . . . . .	483 ff
— lycaonicum . . . . .	369 f*	— rhagadioloides . . . . .	483 ff* m
— orientale . . . . .	365 f*	Hemicarpha . . . . .	473 ff
— pendikum . . . . .	361 f* m	Hibiscus fritzschiae . . . . .	20
— polyactis . . . . .	373 f*	Hierochloë . . . . .	129 ff
— sulphureum . . . . .	362 m	— alpina ssp. alpina . . . . .	156 ff* m
— tureicum . . . . .	364 f*	— — ssp. <i>orthantha</i> . . . . .	161 ff* m
— zoharii . . . . .	371 f*	— australis . . . . .	134 ff* m
Cajanus cajan . . . . .	443*	— hirta ssp. <i>aretiæ</i> . . . . .	150 ff* m
Calliandra haematocephala . . . . .	443*	— — ssp. hirta . . . . .	146 ff* m
Cardamine graeca . . . . .	415*	— odorata ssp. <i>baltica</i> . . . . .	141 ff* m
— hirsuta . . . . .	415 f*	— — ssp. <i>odorata</i> . . . . .	136 ff* m
Carex . . . . .	319 ff	— repens . . . . .	154 ff* m
— capitata . . . . .	323 f m	Homalothecium lutescens . . . . .	222*
— curta . . . . .	324 ff m	— sericeum . . . . .	222*
— — var. <i>robustior</i> . . . . .	326	Iberis sprunieri . . . . .	416*
— macloviana . . . . .	322 f m	Iris foetidissima . . . . .	377*
— magellanica . . . . .	326 ff m	Isolepis . . . . .	479 f
— maritima . . . . .	321 f m	— <i>kernii</i> . . . . .	479
— microglochin . . . . .	328 ff m	— <i>microcephala</i> . . . . .	480
Centaurea arguta . . . . .	377 f*	— <i>rehmannii</i> . . . . .	479
— taganensis . . . . .	379*	Isothecium myurum . . . . .	219*
— webbiana . . . . .	379	Jatropha curcas . . . . .	54*
Ceratophyllum demersum . . . . .	335 ff m	Juncus . . . . .	1 ff, 179 ff, 311 ff, 435 ff
Cerbera thevetia . . . . .	448*	— arcticus . . . . .	311 ff*
Chrysanthemum coronarium . . . . .	380*	— — ×filiformis . . . . .	316*
Cliffortia . . . . .	292 ff*	— balticus . . . . .	315*
Cneorum tricoccon . . . . .	254*	— — ×filiformis . . . . .	316*
Codiaeum sp. . . . .	56*	— biglumis . . . . .	1 f*
Crepis canariensis . . . . .	381*	— capitatus . . . . .	8*
Croton . . . . .	56*	— castaneus . . . . .	2 f*
Cymbalaria aequitriloba . . . . .	262*	— compressus . . . . .	435 f*
Cyperaceae tribus Cyperaceae . . . . .	280 ff	— conglomeratus . . . . .	181*
— tribus Scirpeae . . . . .	280 ff	— — ×effusus . . . . .	184 f*
Cyperus . . . . .	474 f*	— effusus . . . . .	179 f*
— nudicaulis . . . . .	282*	— — ×inflexus . . . . .	183 f*
Dalechampia roezliana . . . . .	42 f*, 58*	— filiformis . . . . .	311*
Dianthus <b>Stamatiidae</b> . . . . .	77 f*	— gerardi . . . . .	439 f
Dichostemma sp. . . . .	60*	— — ssp. <i>atrofuscus</i> . . . . .	441*
Diplocyathium sp. . . . .	61*	— — ssp. <i>gerardi</i> . . . . .	439 f*
Doxantha unguis-cati . . . . .	448*	— inflexus . . . . .	181 f*
Echium giganteum . . . . .	377*	— squarrosum . . . . .	435*
— leucocephalum . . . . .	377*	— stygius . . . . .	2*
— parviflorum . . . . .	252*	— tenuis . . . . .	441*
— simplex . . . . .	377*	— triglumis . . . . .	1*
— strictum . . . . .	377*	Kigelia pinnata . . . . .	448*
— — ssp. <i>exasperatum</i> . . . . .	377*	Laminaria digitata . . . . .	277 ff*
Eranthemum laxiflorum . . . . .	443*	Launaea cervicornis . . . . .	251*
Erismanthus sp. . . . .	56*	Lavatera cretica . . . . .	259*
Euphorbia . . . . .	46 f	— maritima . . . . .	259*
— dracunculoides . . . . .	46 f*	Lemna trisulca . . . . .	335 ff m
— hirta . . . . .	46 f*	Limnalsine . . . . .	187 f
— nivulia . . . . .	46 f*	— diffusa . . . . .	188 ff* m
— rothiana . . . . .	48*	Linaria pelisseriana . . . . .	262*
Euphorbiaceae . . . . .	39 ff*	Linum bienne . . . . .	257*
Euryhynchium striatum . . . . .	222*	Lipocarpha . . . . .	473 ff*
Fritillaria erhardii . . . . .	66 ff*	— chinensis . . . . .	473*
Fumana ericooides . . . . .	254*	— <i>isolepis</i> . . . . .	476 f*
Galaxia obscura . . . . .	83 f*	Lithospermum <b>Goulandriorum</b> . . . . .	355 ff*

<i>Lotus emeroides</i>	382*	<i>Poinsettia</i>	48 f
— <i>tetraphyllus</i>	254*	— <i>geniculata</i>	48*
<i>Mariscus</i>	475*	— <i>pulcherrima</i>	50*
— <i>paradoxus</i>	477 ff*	<i>Polykarpon polycarpoides</i>	252*
<i>Marrubium alternidens</i>	495	<i>Polygala rupestris</i>	260*
<i>Maxia</i>	187 f, 198 ff	<i>Polytrichum commune</i>	224*
— <i>howellii</i>	198 ff* m	— <i>formosum</i>	224*
<i>Melica minuta</i>	260*, 276 f	— <i>juniperinum</i>	224 f*
<i>Merendera filifolia</i>	257*	— <i>longisetum</i>	225*
<i>Micromeria filiformis</i>	257*	— <i>piliferum</i>	225*
<i>Mnium affine</i>	215*	<i>Potentilla Goulandrii</i>	75 f*
— <i>cinctidioides</i>	215*	<i>Pteronia anisata</i>	10 ff*
— <i>cuspidatum</i>	215*	— <i>elata</i>	9 f*
— <i>horneum</i>	215 f*	<i>Ranunculus macrophyllus</i>	260*
— <i>seligeri</i>	217*	<i>Reseda alba</i>	260*
— <i>undulatum</i>	217*	<i>Rhytidadelphus loreus</i>	222 f*
<i>Monadenium laeve</i>	46*	— <i>squarrosum</i>	223*
— <i>schubei</i>	46*	— <i>triquetus</i>	223*
<i>Monsonia parvifolia</i>	208 ff* m	<i>Ricotia cretica</i>	416
<i>Montiastrum</i>	87 ff	<i>Rubia peregrina</i>	262*
— <i>bostockii</i>	92 ff*, 102 ff*	<i>Rumex pulcher</i>	260*
— <i>dichotomum</i>	95 ff* m, 104 ff*	<i>Ruta angustifolia</i>	262*
— <i>lineare</i>	98 ff* m, 103 ff*	— <i>oreojasme</i>	382*
— <i>vassilievii</i>	90 ff*, 102 ff	<i>Salvia verbenaca</i>	257*
<i>Myoporum</i>	458 ff	<i>Samanea saman</i>	443*
— <i>deserti</i>	466 f*	<i>Schoenoplectus</i>	288 f
<i>Myriophyllum spicatum</i>	335 ff m	— <i>brachyceras</i>	290
<i>Nigella arvensis</i> ssp. <i>arvensis</i>	231 ff m	— <i>confusus</i>	290
<i>Oftia</i>	451 ff, 458 ff	— <i>inelinatus</i>	290
— <i>africana</i>	456 ff* m, 464 f*	— <i>jacobi</i>	290
— <i>glabra</i>	454 f*	— <i>lateriflorus</i>	290
— <i>revoluta</i>	454 f*	— <i>microglumis</i>	287 f*
<i>Ononis reclinata</i>	255*	— <i>rogersii</i>	290
<i>Onosma psammophilum</i>	79 f*	— <i>roylei</i>	290
<i>Ophrys fusca</i>	259*	— <i>subulatus</i>	290
<i>Othonna hederifolia</i>	14 f*	<i>Scoparia dulcis</i>	237 ff*
<i>Oxycaryum</i>	281 f	<i>Sideritis gomerae</i>	382*
— <i>cubense</i>	282 f*	<i>Silene nocturna</i>	252*
<i>Pancreatum canariense</i>	376*	— <i>pseudatocion</i>	252*
<i>Parietaria diffusa</i>	262*	<i>Spathodea campanulata</i>	447*
— <i>Iusitanica</i> ssp. <i>Iusitanica</i>	262*	<i>Stipa capensis</i>	260*
<i>Parmelia subgen. Xanthoparmelia</i>	343 ff	<i>Stipagrostis papposa</i>	17 ff
— <i>almhornii</i>	345 f*	— <i>uniplumis</i>	17 ff
— <i>atroventralis</i>	346*	— <i>vexillifera</i>	16 f*
— <i>diadeta</i>	346 f*	<i>Synadenium grantii</i>	44 f*
— <i>dichromatica</i>	348*	<i>Tapeinia</i>	82 ff
— <i>encrustans</i>	348 f*	— <i>obscura</i>	85 f* m
— <i>frondosa</i>	349*	— <i>pumila</i>	82 f
— <i>heterodoxa</i>	349*	<i>Taraxacum</i>	497 ff
— <i>hyporhytida</i>	349 f*	— <i>baeckiforme</i>	499 f*
— <i>lecanorica</i>	351*	<i>Tecomaria capensis</i>	448*
— <i>neoecongensis</i>	351 f*	<i>Teedia</i>	458 ff
— <i>quintaria</i>	353*	— <i>lucida</i>	458 ff*, 468 f*
— <i>subdomokosii</i>	353 f*	— <i>sp.</i>	458 ff*
<i>Pedilanthus tithymaloides</i>	50 f*	<i>Thuidium tamariscinum</i>	217 f*
<i>Phylanthus niruri</i>	56*	<i>Thunbergia alata</i>	443*
<i>Plantago macrorrhiza</i>	259*, 267	— <i>erecta</i>	443*
<i>Platystemma violoides</i>	25 ff*	— <i>fragans</i>	446*
<i>Pleurozium schreberi</i>	222*	<i>Todoroa aurea</i>	382*
<i>Pogonatum aloides</i>	223*	<i>Tribulus echinops</i>	21 f*
— <i>urnigerum</i>	223 f*	<i>Tulipa cretica</i>	66 f*

## VIII

<i>Valerianella coronata</i> .....	263*	<i>Vieraea laevigata</i> .....	381*
— <i>eriocarpa</i> .....	263*	<i>Viola arborescens</i> .....	263*
— <i>microcarpa</i> .....	263*	<i>Vulpia membranacea</i> .....	260*
<i>Vallaris solanacea</i> .....	448*		