

BOTANISKA NOTISER

1970, VOL. 123

ISSUED BY
LUNDS BOTANISKA FÖRENING

EDITED BY
INGEMAR BJÖRKQVIST

Tidskriften grundad 1839

DISTRIBUTOR: C. W. K. GLEERUP, LUND, SWEDEN

Distribution Dates

- Fasc. 1, pp. 1—212, on 10 April 1970
Fasc. 2, pp. 213—362, on 22 June 1970
Fasc. 3, pp. 363—446, on 16 Oct. 1970
Fasc. 4, pp. 447—576, on 30 Dec. 1970

Contents (Innehållsförteckning)

| | |
|--|-----|
| ANDERSSON, F.: Ecological Studies in a Scanian Woodland and Meadow Area, Southern Sweden II. Plant Biomass, Primary Production and Turnover of Organic Matter | 8 |
| BHATNAGAR, S. P. and GUPTA, M.: Morphology and Embryology of <i>Mimusops elengi</i> L. | 455 |
| BOTHMER, R. VON: Studies in the Aegean Flora XV. Chromosome Numbers in Labiatae | 52 |
| — Cytological Studies in <i>Allium</i> I. Chromosome Numbers and Morphology in <i>Allium</i> Sect. <i>Allium</i> from Greece | 518 |
| BRAMWELL, D.: A Revision of the Genus <i>Parolinia</i> Webb (Cruciferae) in the Canary Islands | 394 |
| DAHLGREN, R.: <i>Wiborgia apterophora</i> R. Dahlgr., a New Species of Leguminosae from the Cape Province | 112 |
| — Current Topics, Parallelism, Convergence, and Analogy in Some South African Genera of Leguminosae | 551 |
| DIXON, P. S. and IRVINE, L. M.: Miscellaneous Notes on Algal Taxonomy and Nomenclature III | 474 |
| EKBERG, L.: Studies in the Genus <i>Allium</i> III. Wind Dispersal of <i>Allium</i> Bulbs .. | 115 |
| ELIASSON, U.: Studies in Galápagos Plants VIII. Chromosome Numbers of Some Endemic Species | 149 |
| — Studies in Galápagos Plants IX. New Taxonomical and Distributional Records | 346 |
| ENGSTRAND, L.: Studies in the Aegean Flora XVIII. Notes and Chromosome Numbers in Aegean Umbelliferae | 384 |
| — The European Species of <i>Scaligeria</i> (Umbelliferae) | 505 |
| FLENSBURG, T. and MALMER, N.: Studies on Mire Vegetation in the Archaean Area of South-Western Götaland (South Sweden) IV. Benthic Algae and Their Distribution on the Åkhull Mire | 269 |
| GUSTAFSSON, M.: Studies in the Aegean Flora XVII. Variation and Distribution of <i>Atriplex recurva</i> D'Urv. (Chenopodiaceae) | 371 |
| HEDBERG, O.: The Genus <i>Zaluzianskya</i> F. W. Schmidt (Scrophulariaceae) Found in Tropical East Africa | 512 |
| KNABEN, G. and HYLANDER, N.: On the Typification of <i>Papaver radicatum</i> Rottb. and its Nomenclatural Consequences | 338 |
| LINDGREN, L.: <i>Stellaria nemorum</i> ssp. <i>glochidosperma</i> Found in Småland, South Sweden (<i>Stellaria nemorum</i> ssp. <i>glochidosperma</i> funnen i Småland) | 200 |
| — Beech Forest Vegetation in Sweden — a Survey | 401 |

IV

| | |
|---|-----|
| LYE, K. A. and HAINES, R. H.: Studies in African Cyperaceae I. A New Species of <i>Scirpus</i> (Cyperaceae) from Mt. Elgon | 430 |
| MOE, D.: The Post-Glacial Immigration of <i>Picea abies</i> into Fennoscandia | 61 |
| NIHLGÅRD, B.: Vegetation Types of Planted Spruce Forests in Scania, Southern Sweden | 310 |
| NILSSON, J.: Notes on the Biomass and Productivity of Below-Ground Organs of a South-Swedish Hay-Meadow | 183 |
| NILSSON, Ö.: Studies in <i>Montia</i> L., <i>Claytonia</i> L. and Allied Genera IV. The Genus <i>Crunocallis</i> Rydb. | 119 |
| NORDENSTAM, B.: Studies in South African Liliaceae III. The Genus <i>Rhadamanthus</i> | 155 |
| — Notes on South African Iridaceae: <i>Lapeirousia</i> and <i>Babiana</i> | 433 |
| OREDSSON, A.: Drawings of Scandinavian Plants 33—36. <i>Rubus</i> L. Subgen. <i>Rubus</i> | 1 |
| — Ditto 37—40. <i>Rubus</i> L. Subgen. <i>Rubus</i> | 213 |
| — Ditto 41—44. <i>Rubus</i> L. Subgen. <i>Rubus</i> | 363 |
| — Ditto 17—44. Key to <i>Rubus</i> L. Subgen. <i>Rubus</i> | 447 |
| RAVANKO, O.: Observations on the Taxonomy and Ecology of the Epilithic Crustaceous Brown Algae in the SW Archipelago of Finland (Seili Islands) | 220 |
| SNOGERUP, S.: Studies in the Genus <i>Juncus</i> IV. The Typification of <i>Juncus conglomeratus</i> L. | 425 |
| SONESSON, M.: Studies on Mire Vegetation in the Torneträsk Area, Northern Sweden IV. Some Habitat Conditions of the Poor Mires | 67 |
| THULIN, M.: Chromosome Numbers of Some Vascular Plants from East Africa .. | 488 |
| WEIMARCK, G.: <i>Hierochloë alpina</i> (Gramineae) $2n=66$ in Northern Finland | 201 |
| — Spontaneous and Induced Variation in Some Chemical Leaf Constituents in <i>Hierochloë</i> (Gramineae) | 231 |
| — Apomixis and Sexuality in <i>Hierochloë alpina</i> (Gramineae) from Finland and Greenland and in <i>H. monticola</i> from Greenland | 495 |
| WENDELBO, P.: New and Noteworthy Species of Primulaceae from the 'Flora Iranica'-Area | 300 |
| Berichtigung | 199 |
| Lunds Botaniska Förening (The Lund Botanical Society) | |
| Från Lunds Botaniska Förenings förhandlingar 1969 | 195 |
| Lunds Botaniska Förening 1970 | 574 |
| Reviews of Botanical Literature | |
| BJÖRKQVIST, I.: Studies in <i>Alisma</i> I—II. (By the Editor) | 570 |
| BOLIN, L. and NILSSON, Ö.: Fotofloran. (By L. LINDGREN) | 210 |
| BRIGGS, G. E.: Movements of Water in Plants (By L. PERSSON) | 204 |
| BUTENKO, R. G.: Plant Tissue Culture and Plant Morphogenesis. (By L. O. BJÖRN) .. | 208 |
| COLE, A. J. (ed.): Numerical Taxonomy. (By G. WEIMARCK) | 444 |
| GOULANDRIS, N. A., GOULIMIS, C. N. and STEARN, W. T.: Wild Flowers of Greece. (By A. STRID) | 569 |
| HEATH, O. V. S.: The Physiological Aspects of Photosynthesis. (By L. O. BJÖRN) .. | 206 |

| | |
|--|----------|
| HEGNAUER, R.: Chemotaxonomie der Pflanzen. (By U. OLSSON) | 203 |
| LEVRING, T., HOPPE, H. A. and SCHMID, O. J.: Marine Algae. A Survey of Research and Utilization. (By T. VON WACHENFELDT) | 211 |
| RICHTER, G.: Die Stoffwechselfysiologie der Pflanzen. (By L. O. BJÖRN) | 207 |
| ROBERTSON, R. N.: Protons, Electrons, Phosphorylation, and Active Transport. (By S. PETERSSON) | 205 |
| ROUND, F. E.: Introduction to the Lower Plants. (By G. WEIMARCK) | 209 |
| SCHMITHÜSEN, J.: Allgemeine Vegetationsgeographie. (By N. MALMER) | 360 |
| SINGH, A.: Plant Physiology. (By L. O. BJÖRN) | 361 |
| SWANSON, C. P., MERZ, T. and YOUNG, W. J.: Zytogenetik. (By A. STRID) | 358 |
| STILES, W. and COCKING, E. C.: An Introduction to the Principles of Plant Physiology. (By L. O. BJÖRN) | 359 |
| TYLER, G.: Studies in the Ecology of Baltic Sea-Shore Meadows II. (By the Editor) | 572 |
| Announcements | 199, 573 |

Index (Register)

The index covers only taxa which have been treated in more detail. New taxa and new combinations are printed in boldface. * indicates illustration(s) (also of chromosomes) and m refers to a distribution map.

Registret upptager endast taxa som i något avseende blivit mera utförligt behandlade. Nya taxa och nya kombinationer betecknas med fetstil. * anger avbildning eller avbildningar (även av kromosomer) och m betecknar utbredningskarta.

| | | | |
|--|----------------|---|-----------|
| Allium | 115 ff, 518 ff | — recurva | 371 ff* m |
| — ampeloprasum | 533 f* | — tatarica | 371 ff* |
| — bimetrale | 533* m | Babiana sect. Acaste | 440 |
| — bourgaei | 533* m | — angustifolia | 440 f |
| — descendens | 535 f* | — horizontalis | 439 f |
| — sphaerocephalum | 534 f* | — lewisiana | 436 ff* |
| — stocksianum | 117 f* | — multiflora | 442 f* |
| Amaracus tournefortii | 52* | — nana var. angustifolia | 442 |
| Ammi majus | 385 | — — var. maculata | 442 |
| Anagallis serpens ssp. meyeri- johannis | 489* | — parviflora | 443 |
| Androsace podlechii | 302* | — pulchra | 440 f |
| Anthemis tigreensis | 489* | Borbonia | 556 f* |
| Anthriscus caucalis | 385 | Bowlesia palmata | 352 f |
| Apium graveolens | 385 | Calamintha exigua | 52 f* |
| Artemisia squamata | 385 | Carum multiflorum ssp. multiflorum | 385 f* |
| Aspalathus | 554 ff, 563 ff | Coryza subscaposa | 489* |
| — acanthoclada | 556* | Coridothymus capitatus | 53* |
| — aculeata | 564* | Corydalis mildbraedii | 489* |
| — acuminata | 566* | Cosmarium pseudoexiguum | 280* |
| — alpestris | 566* | Crassula granvikii | 489* |
| — chenopoda | 564* | Crunocallis | 121 ff |
| — ericifolia | 566* | — calcicola | 138 ff* m |
| — glabrata | 566* | — chamissoi | 124 ff* m |
| — heterophylla | 566* | Cynorkis anacamptoides | 490* |
| — lanata | 556 f* | Dasya baillouviana | 482 f* |
| — lanifera | 556 f* | — pedicellata | 480 ff |
| — radiata | 554 f* | Daucus carota ssp. drepanensis ... | 386* |
| — sericea | 554 f* | — guttatus ssp. guttatus | 387* |
| — spinosissima | 565* | — involucreatus | 387* |
| Atriplex halimus | 373 ff* m | Dierama pendulum | 490* |

| | | | |
|---|------------------|--|-------------------------------|
| <i>Dionysia esfandiarii</i> | 302 f* | <i>Orlaya kochii</i> | 390 |
| — freitagii | 303 f* | <i>Papaver radicum</i> | 338 ff* |
| — termiana | 306* | <i>Parolinia</i> | 394 ff |
| <i>Euchlora</i> | 553 f | — intermedia | 398 f* m |
| Fabaceae tribus Borbonieae | 552 ff | — ornata | 395 f* m |
| <i>Fagus sylvatica</i> | 401 ff m | — schyzogynoides | 396* m |
| — communis | 387 | <i>Pennisetum schimperii</i> | 490* |
| <i>Ferula chiliantha</i> | 387* | <i>Pernettya howellii</i> | 353 |
| <i>Grateloupia</i> | 475 ff | <i>Petroderma maculiforme</i> | 220 ff |
| <i>Halymenia latifolia</i> | 477 f* | <i>Picea abies</i> | 61 ff m |
| <i>Haplocarpha rueppellii</i> | 489* | <i>Pimpinella peregrina</i> | 390 |
| <i>Heribaudiella fluviatilis</i> | 228 | <i>Pleuroptalum darwinii</i> | 150* |
| <i>Hierochloë</i> | 231 ff, 495 ff | <i>Pleurotaenium tridentulum</i> | 286* |
| — alpina | 201*, 497 ff* | <i>Polysiphonia</i> | 478 f |
| — monticola | 501 | Portulacaceae subfam. | |
| <i>Juncus conglomeratus</i> | 425 ff* | Montioideae | 119 ff |
| — conglomeratus × effusus | 428* | <i>Prasium majus</i> | 54 f* |
| — effusus | 428 f* | <i>Primula algida</i> | 306 f |
| <i>Koeleria capensis</i> | 490* | — dueckelmannii | 307* m |
| <i>Lagoccia cuminoides</i> | 388 | — gaubaeana | 307 f m |
| <i>Lapeirousia lewisiana</i> | 434 f* | — kaufmanniana | 309* |
| <i>Lavandula stoechas</i> | 53 | <i>Prunella vulgaris</i> | 56* |
| <i>Lebeckia</i> | 558 ff | <i>Pseudorlaya pumila</i> | 390 |
| — cinerea | 558 f* | <i>Rafnia</i> | 563* |
| — halenbergensis | 560* | <i>Rhadamanthus</i> | 153 ff |
| — linearifolia | 558 f* | — albiflorus | 177 f* m |
| — meyeriana | 560* | — arenicola | 166 ff* m |
| — plukenetiana | 560* | — convallarioides | 159 ff* m |
| <i>Lecocarpus foliosus</i> | 153* | — cyanelloides | 179 f* m |
| <i>Lithoderma fatiscens</i> | 227 f* | — fasciatus | 174 f* m |
| — roseningii | 220 ff | — montanus | 162 ff* m |
| — subextensum | 220 ff | — platyphyllus | 172 ff* m |
| <i>Lobelia aberdarica</i> | 489* | — urantherus | 164 ff* m |
| — cf. duriprati | 489* | <i>Rubus</i> | 1 ff, 213 ff, 363 ff, 447 ff* |
| <i>Lotononis</i> | 553 f | — armeniacus | 1 ff* |
| — azurea | 553* | — bellardii | 363 ff* |
| — peduncularis | 555* | — caesius | 369 f* |
| — serpens | 555* m | — corylifolius agg. | 368 f* |
| <i>Lysimachia ruhmeriana</i> | 489* | — fuscus | 217 ff* |
| <i>Majorana onites</i> | 53* | — gothicus | 366* |
| <i>Malabaila involucreta</i> | 388* | — hartmani | 363* |
| <i>Marrubium vulgare</i> | 54* | — lindebergii | 1* |
| <i>Mentha pulegium</i> | 54* | — radula | 213 f* |
| <i>Miroscladum minutum</i> | 388* | — taeniarum | 216 f* |
| <i>Mimusops elengi</i> | 455 ff* | — thyrsanthus | 6* |
| <i>Nama dichotomum</i> | 353 | — vestervicensis | 6 f* |
| <i>Nolana galapagensis</i> ... | 150 f*, 351 ff m | — vestitus | 213* |
| <i>Origanum heracleoticum</i> | 54* | — wahlbergii | 366* |

VIII

| | | | |
|--|---------------|--------------------------------------|-----------|
| <i>Salvia pomifera</i> ssp. <i>calycina</i> .. | 56 f* m | <i>Sebaea brachyhylla</i> | 489* |
| — — ssp. <i>pomifera</i> | 57* m | <i>Senecio</i> sp. | 489* |
| — <i>verbenacea</i> ssp. <i>multifida</i> | 58* | <i>Sida rupo</i> | 350 ff* |
| — <i>viridis</i> | 58 f* | <i>Smyrnum rotundifolium</i> | 391* |
| <i>Satureja thymbra</i> | 59* | <i>Sporobolus olivaceus</i> | 490* |
| <i>Scalesia</i> | 152 | <i>Staurostrum longispinum</i> | 287* |
| — <i>affinis</i> | 152* | <i>Stellaria nemorum</i> ssp. | |
| — <i>crockeri</i> | 152 f* | <i>glochidosperma</i> | 200 |
| — <i>hopkinsii</i> | 153* | <i>Swertia crassiuscula</i> | 489* |
| — <i>microcephala</i> | 153* | <i>Torilis leptophylla</i> | 391 f* |
| <i>Scaligeria</i> | 505 ff | — <i>nodosa</i> | 393* |
| — <i>cretica</i> | 390 f* | <i>Urocarpidium insulare</i> | 348 ff* |
| — <i>halophila</i> | 391*, 508 f m | <i>Wiborgia apterophora</i> | 112 ff* |
| — moreana | 509 f* m | <i>Xiphopteris serrulata</i> | 346 f* |
| — <i>napiformis</i> | 506 f m | <i>Xyris capensis</i> | 490* |
| <i>Scandix australis</i> ssp. <i>australis</i> ... | 391* | <i>Zaluzianskya</i> | 512 ff |
| <i>Scinaia</i> | 474 f | — elgonensis | 512 ff* m |
| — <i>forcellata</i> | 475 | — <i>gilioides</i> | 513 m |
| <i>Scirpus graminoides</i> | 430 f* | — <i>pusilla</i> | 513 m |